

SURAT KETERANGAN

Nomor: 036/UNUSA/Adm-LPPM/I/2020

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 15 Januari 2020.

Judul : Relationships Between Lead Contaminated Seafood Consumption And Blood Pressure Among Fisherman Communities At The Makassar Coastal Areas Indonesia

Penulis : Anwar Mallongi, Agus Bintara Birawida, Apollo, Abdul Muhith, Jastal

Identitas : Indian Journal of Physiotherapy and Occupational Therapy, Vol 12, No.4 October 2018

No. Pemeriksaan : 2020.01.17.26

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 16%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya

Surabaya, 17 Januari 2020

Ketua LPPM,



Dr. Istas Pratomo, S.T., M.T.

NPP. 16081074

Paper 4

by Abdul Muhith 4

Submission date: 15-Jan-2020 01:57PM (UTC+0700)

Submission ID: 1242145449

File name: Volume_12,_Number_4_IJPOT_IPOT_IJPOT_October_December_2018..pdf (839.52K)

Word count: 5282

Character count: 31890

ISSN P - 0973-5666
ISSN E - 0973-5674

Volume 12

Number 4

October-December 2018

Indian Journal of Physiotherapy and Occupational Therapy

An International Journal



website: www.ijpot.com

Indian Journal of Physiotherapy and Occupational Therapy

Editor-in-Chief

Archna Sharma

Ex-Head Dept. of Physiotherapy, G. M. Modi Hospital, Saket, New Delhi - 110 017

Email : editor.ijpot@gmail.com

Sub-Editor

Kavita Behal

MPT (Ortho)

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Vikram Mohan** (*Lecturer*) Universiti Teknologi MARA, Malaysia
2. **Angusamy Ramadurai** (*Principal*) Nyangabgwe Referral Hospital, Botswana
3. **Faizan Zaffar Kashoo** (*Lecturer*) College Applied Medical Sciences, Al-Majma'ah University, Kingdom of Saudi Arabia
4. **Amr Almaz Abdel-aziem** (*Assistant Professor*) of Biomechanics, Faculty of Physical Therapy, Cairo University, Egypt
5. **Abhilash Babu Surabhi** (*Physiotherapist*) Long Sault, Ontario, Canada
6. **Avanianban Chakkarapani** (*Senior Lecturer*) Quest International University Perak, IPOH, Malaysia
7. **Manobhram Nellutia** (*Safety Advisor*) Fiosa-Miosa Safety Alliance of BC, Chilliwack, British Columbia
8. **Jaya Shanker Tedla** (*Assistant Professor*) College of Applied Medical Sciences, Saudi Arabia
9. **Stanley John Winsor** (*PhD Candidate*) at University of Otago, New Zealand
10. **Salwa El-Sobkey** (*Associate Professor*) King Saud University, Saudi Arabia
11. **Saleh Aloraibi** (*Associate Professor*) College of Applied Medical Sciences, Saudi Arabia
12. **Rashid M.** (*Faculty-PT Neuro Sciences*) College of Allied Health Sciences, UAE
13. **Mohmad Waseem**, (*Exercise Therapist*) Alberta- CANADA
14. **Muhammad Naveed Babur** (*Principle & Associate Professor*) Isra University, Islamabad, Pakistan
15. **Zbigniew Sliwinski** (*Professor*) Jan Kochanowski University in Kielce
16. **Mohammed Taher Ahmed Omar** (*Assistant Professor*) Cairo University, Giza, Egypt
17. **Ganesan Kathiresan** (*DBC Senior Physiotherapist*) Kuching, Sarawak, Malaysia
18. **Kartik Shah** (*Health Consultant*) for the Yoga Expo, Canada
19. **Shweta Gore** (*Senior Physical Therapist*) Narayan Rehabilitation, Bad Axe, Michigan, USA
20. **Ashokan Arumugam** MPT (*Ortho & Manual Therapy*), PhD, Department of Physical Therapy, College of Applied Medical Sciences, Majmaah University, Kingdom of Saudi Arabia
21. **Veena Raigangar** (*Lecturer*) Dept. of Physiotherapy University of Sharjah, U.A.E
22. **Dave Bhargav** (*Senior Physical Therapist*) Houston, Texas
23. **Dr. Jagatheesan A**
Assistant Professor, Gulf Medical University, Ajman, UAE.

NATIONAL EDITORIAL ADVISORY BOARD

1. **Charu Garg** (*Incharge PT*), Sikanderpur Hospital (MJSMS), Sirsa Haryana, India
2. **Vaibhav Madhukar Kapre** (*Associate Professor*) MGM Institute of Physiotherapy, Aurangabad (Maharashtra)
3. **Amit Vinayak Nagrale** (*Associate Professor*) Maharashtra Institute of Physiotherapy, Latur, Maharashtra
4. **Manu Goyal** (*Principal*), M.M University Mullana, Ambala, Haryana, India
5. **P. Shanmuga Raju** (*Asst. Professor & I/C Head*) Chalmeda AnandRao Institute of Medical Sciences, Karimnagar, Andhra Pradesh
6. **Sudhanshu Pandey** (*Consultant Physical Therapy and Rehabilitation*) Department / Base Hospital, Delhi
7. **Aparna Sarkar** (*Associate Professor*) AIPT, Amity university, Noida
8. **Jasobanta Sethi** (*Professor & Head*) Lovely Professional University, Phagwara, Punjab
9. **Patitapaban Mohanty** (*Assoc. Professor & H.O.D*) SVNIRTAR, Cuttack, Odisha
10. **Suraj Kumar** (*HOD and Lecturer*) Physiotherapy Rural Institute of Medical Sciences & Research, Paramedical Vigyan Mahavidhyalaya Saifai, Etawah, UP
11. **U. Ganapathy Sankar** (*Vice Principal*) SRM College of Occupational Therapy, Kattankulathur, Tamil Nadu
12. **Hemant Juneja** (*Head of Department & Associate Professor*) Amar Jyoti Institute of Physiotherapy, Delhi
13. **Sanjiv Kumar** (*I/C Principal & Professor*) KLEU Institute of Physiotherapy, Belgaum, Karnataka
14. **Shaji John Kachanathu** (*Associate Professor*) Jaipur Physiotherapy College, Rajasthan, India
15. **Narasimman Swaminathan** (*Professor, Course Coordinator and Head*) Father Muller Medical College, Mangalore
16. **Pooja Sharma** (*Assistant Professor*) AIPT, Amity University, Noida
17. **Nilima Bedekar** (*Professor, HOD*) Musculoskeletal Sciences, Sancheti Institute College of Physiotherapy, Pune.
18. **N. Venkatesh** (*Principal and Professor*) Sri Ramachandra university, Chennai
19. **Meenakshi Batra** (*Senior Occupational Therapist*), Pandit Deen Dayal Upadhyaya Institute for The Physically Handicapped, New Delhi
20. **Shovan Saha, T** (*Associate Professor & Head*) Occupational Therapy School of Allied Health Sciences, Manipal University, Manipal, Karnataka
21. **Akshat Pandey**, *Sports Physiotherapist*, Indian Weightlifting Federation / Senior Men and Woman / SAI NSNIS Patiala

Indian Journal of Physiotherapy and Occupational Therapy

NATIONAL EDITORIAL ADVISORY BOARD

23. **Maneesh Arora**, Professor and as Head of Dept, Sardar Bhagwan (P.G.) Institute of Biomedical Sciences, Balawala, Dehradun, UK
24. **Deepak Sharan**, Medical Director and Sole Proprietor, RECOUP Neuromusculoskeletal Rehabilitation Centre, New Delhi
25. **Jayaprakash Jayavelu**, Chief Physiotherapist –Medanta The Medicity, Gurgaon Haryana
26. **Vaibhav Agarwal**, Incharge, Dept of physiotherapy, HIHT, Dehradun,
27. **Shipra Bhatia**, Assistant Professor, AIPT, Amity university, Noida
28. **Jaskirat Kaur**, Assistant Professor, Indian Spinal Injuries Center, New Delhi
29. **Prashant Mukkanavar**, Assistant Professor, S.D.M College of Physiotherapy, Dharwad, Karnataka
30. **Chandan Kumar**, Associate Professor & HOD, Neuro-physiotherapy, Mahatma Gandhi Mission's Institute of Physiotherapy, Aurangabad, Maharashtra
31. **Satish Sharma**, Assistant Professor, I.T.S. Paramedical College Murad Nagar Ghaziabad
32. **Richa**, Assistant Professor, I.T.S. Paramedical College Murad Nagar Ghaziabad
33. **Manisha Uttam**, Research Scholar, Punjabi University, Patiala
34. **Dr. Ashfaq Khan (PT)**, HOD Physiotherapy, Integral University Lucknow U.P.,
35. **Dr. Dibyendunaryan Bid (PT)** Senior Lecturer The Sarvajani College of Physiotherapy Rampura, Surat
36. **Vijayan Gopalakrishna Kurup**, Senior Physiotherapist, Rajagiri Healthcare & Education Trust, Aluva, Kerala

SCIENTIFIC COMMITTEE

1. **Gaurav Shori** (Assistant Professor) I.T.S College of Physiotherapy
2. **Baskaran Chandrasekaran** (Senior Physiotherapist) PSG Hospitals, Coimbatore
3. **Dharam Pandey** (Sr. Consultant & Head of Department) BLK Super Speciality Hospital, New Delhi

SCIENTIFIC COMMITTEE

4. **Jeba Chitra** (Associate Professor) KLEU Institute of Physiotherapy Belgaum, Karnataka
5. **Deepak B. Anap** (Associate Professor) PDVPPF's, College of Physiotherapy, Ahmednagar. (Maharashtra)
6. **Shalini Grover** (Assistant Professor) HOD-FAS, MRIU
7. **Vijay Batra** (Lecturer) ISIC Institute of Rehab. Sciences
8. **Ravinder Narwal** (Lecturer) Himalayan Hospital, HIHT Medical University, Dehradun-UK.
9. **Abraham Samuel Babu** (Assistant Professor) Manipal College of Allied Health Sciences, Manipal
10. **Anu Bansal** (Assistant Professor and Clinical Coordinator) AIPT, Amity University, Noida
11. **Bindya Sharma** (Assistant Professor) Dr. D. Y. Patil College of Physiotherapy, Pune
12. **Dheeraj Lamba**, Associate Professor & Research Coordinator, School of Physiotherapy, Lovely Professional University, Phagwara (India)
13. **Soumya G** (Assistant Professor) (MSRMC)
14. **Nalina Gupta Singh** (Assistant Professor) Physiotherapy, Amar Jyoti Institute of Physiotherapy, University of Delhi
15. **Gayatri Jadav Upadhyay** (Academic Head) Academic Physiotherapist & Consultant PT, RECOUP Neuromusculoskeletal Rehabilitation Centre, Bangalore
16. **Nusrat Hamdani** (Asst. Professor and Consultant) Neurophysiotherapy (Rehabilitation Center, Jamia Hamdard) New Delhi
17. **Ramesh Debur Visweswara** (Assistant Professor) **M.S. Ramaiah Medical College & Hospital, Bangalore**
18. **Nishat Qudus** (Assistant Professor) Jamia Hamdard, New Delhi
19. **Anand Kumar Singh**, Assistant Professor, RP Indraprast Institute of Medical Sciences Karnal, Haryana
20. **Pardeep Pahwa**, Lecturer, Composite Regional Rehabilitation Centre, Sunder-Nagar under NIVH (Ministry of Social Justice & Empowerment, New Delhi)

"Indian Journal of Physiotherapy and Occupational Therapy" An essential indexed peer reviewed journal for all physiotherapists & occupational therapists provides professionals with a forum to discuss today's challenges- identifying the philosophical and conceptual foundations of the practice; sharing innovative evaluation and treatment techniques; learning about and assimilating new methodologies developing in related professions; and communicating information about new practice settings. The journal serves as a valuable tool for helping therapists deal effectively with the challenges of the field. It emphasizes articles and reports that are directly relevant to practice. The journal is now covered by INDEX COPERNICUS, POLAND and covered by many internet databases. The Journal is registered with Registrar of Newspapers for India vide registration number DELENG/2007/20988

Print- ISSN: 0973-5666, Electronic - ISSN: 0973-5674, Frequency: Quarterly (4 issues per volume).

Website: www.ijpot.com

© All Rights reserved The views and opinions expressed are of the authors and not of the Indian Journal of Physiotherapy and Occupational Therapy. The Indian Journal of Physiotherapy and Occupational Therapy does not guarantee directly or indirectly the quality or efficacy of any products or service featured in the advertisement in the journal, which are purely commercial.

Editor

Archna Sharma

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Printed, published and owned by

Archna Sharma

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Published at

Institute of Medico-legal Publications

Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)



Indian Journal of Physiotherapy and Occupational Therapy

www.ijpot.com

CONTENTS

Volume 12, Number 4

October-December 2018

35. Relationships between Lead Contaminated Seafood Consumption and Blood Pressure among Fisherman Communities at the Makassar Coastal Areas, Indonesia 186
Anwar Mallongi, Agus Bintara Birawida, Apollo, Abdul Muhith, Jastal
36. Comparison of Needles Sharpness, Suture Strength, Pain Revocation, Postoperative Scar between Polyvinylidene Flouride and Polypropylene Surgical Suture 192
Warsinggih, Prihantono
37. Recovery of Autologous and Heterologous Serum Plus Nata De Coco Patch at Ruptured Tympanic Membrane in Laying Hen 198
Syahrijuita, Abdul Kadir, Sartini, Marhaen Hardjo, Burhanuddin Bahar
38. Extract of Elicited Kaffir Lime (*Citrus hystrix* DC.) Cells Suspension by *Saccharomyces cerevisiae* H. and Its Citotoxicity Against T47D Cells 202
Dwi Priyanto, Woro Anindito Sri Tunjung, Ari Indriyanto
39. Indicators of Low Back Pain in Dockworkers in Soekarno Hatta Seaport of Makassar 210
Syamsiar S Russeng, Lalu Muhammad Saleh, Masyitha Muis, Asti Hardianti Azis, Devintha Virani, Anwar Mallongi
40. Personal Management and Concept of a Woman Suffering Cervix Cancer 215
Sandu Siyoto
41. Effectiveness of Health Education using Audio Visual with Lectures and Poster with Lectures of Changes in Behavior Selection of Snacks 220
Nurwijayanti, Nur Yeny Hidajaturokhmah, Maulina Nurikasari, Erma Retnaningtyas, Retno Palupi Yonni Siwi, Nia Sari, Muhammad Ali Sodik
42. Working Climate with Fatigue Levels of Timber Industry Workers in Probolinggo 225
Haman Mujib, Muhith Abdul, Lantin Sulistyorini, Mallongi Anwar, Arief F, Asih M.Y, Vira A
43. Family Support as a Determinant Safety Riding Student Behavior in SMKN 2 Kediri 230
Nia Sari, Amarin Yudhana, Chandra Wahyuni, Aprin Rusmawati, Indasah, Muhammad Ali Sodik
44. Basic Life Support Training Effect (BLS) in the Lay of Knowledge and Behavior Management Cardiac Arrest in RT 14 RW 03 Blimbing Sub District Dongko Trenggalek 235
Novita Ana Anggraini, Rahmania Ambarika, Alfian Fawzi, Suci Anggraeni, Agusta Dian Ellina, Shanty Natalia
45. Histopathologic Analysis of Contralateral Testes in Wistar Rats given Exogen Gluthathion after Testicular Torsion-Detorsion 240
Nita Mariana, Moch. Hatta, Suriani Bahrin, Ridwan, Fatoni Lubis, Faiqah Umar

46. Health Counseling Effect Towards Knowledge Levels of Community in Obtaining Clean Water Supply	246
<i>Abdul Ganing</i>	
47. Carbapenemase Production Screening on Gram-Negative Bacteria Isolate from Waste Material of Mamuju Regency Regional General Hospitals	251
<i>Hasir</i>	
14 48. Effect of Giving Steamed Yellow Pumpkin to Decrease Blood Glucose Level in Type II Diabetes Mellitus Patients in Work Area of Health Center of Mamuju District	257
<i>Imelda Appulembang</i>	
49. The Effect of Tembelekan Leaves Extract (<i>Lantana Camara</i> L) as Electrical Rechargeable of Mosquitoes Repellent to Eliminate <i>Aedes Aegypti</i>	261
<i>Zrimurti Mappau, Abdul Ganing</i>	
50. The Healing of Perforated Tympanic Membrane with Nata De Coco Patch	267
<i>Syahrijuita, Abdul Kadir, Sartini, Marhaen Hardjo, Burhanuddin Bahar</i>	
51. Case Control Study: Influence Factor of Exclusive Breastfeeding Primipara in Rural Pinrang Regency	271
<i>A Ummu Salmah, Sri wanty, Ansariadi, St. Fatimah H</i>	
52. Dental Health Child with Retardation Mental and Parents Behavior	278
<i>Muhammad Ali Sodik, Nita Dwi Astikasari, Intan fazrin, Arina Chusnatayaini, Yuly Peristiowati, Nurwijayanti</i>	

Relationships between Lead Contaminated Seafood Consumption and Blood Pressure among Fisherman Communities at the Makassar Coastal Areas, Indonesia

Anwar Mallongi¹, Agus Bintara Birawida¹, Apollo², Abdul Muhith³, Jastal⁴

¹Environmental Health Department, Faculty of Public Health, Hasanuddin University, Makassar South Sulawesi, Indonesia, ²Politeknik Negeri Ujung Pandang, Sulawesi Selatan, ³Associate Professor in Nursing Department of STIKes Majapahit, ⁴Health Research and Development Unit, Banjarnegara, Indonesia

ABSTRACT

Background: Lead contaminated coastal areas have been widely studied in many cities both in high-income countries and in some developing countries. However, the related health disturbance outcomes due to the lead seafood consumption have not been well documented particularly in low and middle income countries such as in Indonesia particularly in Makassar city where no data available. This research aimed to investigate the relationships between lead seafood consumption, blood lead level (BLL), and blood pressure (BP) and the hypertension in the community-based study site of coastal areas Makassar city, Indonesia.

Method: The number of respondents within this study was 35 adults male that randomly selected, and voluntary base. All respondent sign an inform consent without any force before involved in the research. Information of education, family income, lifestyle, occupational, dietary, smoking habit was gathered by administered household questionnaire interview. Then, systolic blood pressure (SBP) and diastolic blood pressure (DBP) were measured as well as the BLL were measured by inductively coupled plasma mass spectrometry technique. In order to assess the relationships between BLL with SBP and DBP, with the hypertension possibility, multiple linear and logistic regressions were applied.

Results: Pb levels in blood averaging of 27.6 µg/dL with standard deviation 17.56 whereas the minimum value 2 and maximum value 89. In addition, the mean of systolic blood pressure 144.6, standard deviation was 17.56, minimum value was 89 mmHg and value maximum 123 mmHg, mean diastolic blood pressure 101.2, standard deviation 12.37, and the minimum value 54 mmHg and a maximum value of 154 mmHg. Chi square test resulted that there is a relationship between blood pressure and the level of lead in the blood, with p value was 0.01 significant.

Conclusion: The Blood lead level was positively associated with diastolic blood pressure and with the odds for hypertension in adults aged 40 or older. It is necessary to have a monitoring of lead exposure among the fishery communities along the Makassar coastal area.

Keywords: Contaminated coastal, seafood consumption, blood pressure, systolic blood pressure, diastolic and blood pressure

INTRODUCTION

Lead contaminated coastal areas have been widely studied in many cities both in high-income countries and in some developing countries. However, the related health disturbance outcomes due to the lead seafood consumption have not been well documented particularly in low- and middle-income countries such as in Indonesia

particularly in Makassar city. Specifically, data relate to the Blood Lead Level (BLL) is scarcely available for those people who are living along the coastal area of the city. Lead exposure from air, water and soil among adult both women and man are occurred daily and potentially generate hypertension as a results of the increase of blood pressure which may generated from the frequently lead exposure. This research aimed to investigate the

relationships between lead seafood consumption, BLL, and blood pressure (BP) and the hypertension in the community-based study site of coastal areas Makassar city, Indonesia.

A long period of lead exposure and accumulation in the body may generate to effect of serious health. The effects such as; the increased blood pressure, which is associated with the cardiovascular diseases which has been linked to lead exposure^{1,2}. Then hypertensive effects of lead have been widely reported in workers exposed to high levels of the metal and by experimental studies in which animals were exposed to long-term high doses of lead. Under occupational conditions, the development of hypertension has been implicated as a possible consequence of the nephropathy caused by lead exposure. Studies have concluded that lead exposure is a risk factor for raised blood pressure and hypertension even in the general population¹⁻⁴.

Specifically, data relate to the blood lead level (BLL) is scarcely available for those people who are living along the coastal area of the Makassar city. The raised of blood pressure prevalence among adults aged average of 25 years or older was about 35% in 2010.⁵ Lead exposure from air, water, oil and sediment among adult both women and man are occurred daily and potentially generate hypertension as a results of the increase of blood pressure which may generated from the daily lead exposure. Having those described data, this study aimed to examine the relationship of aquatic food consumption with the blood pressure among communities along the Makassar coastal area who are living for more than 25 years on the site.⁶

MATERIALS AND METHOD

The population included adults aged of 40 years or older residents and children in an urban coastal Barombong area of South Sulawesi, Indonesia. For the characteristics respondents, the information on sex, age (years), education level, income class, working period and working duration was obtained from each respondent by the orally administered household interviews⁷. All participants were randomly selected and took part in a household interview. In addition, anthropometric measurement, blood pressure measurements and laboratory tests were conducted individually. Data of 35 interviewed respondents who had performed blood tests and had blood pressure measurements were taken for

analysis purpose to the laboratory. Furthermore, the body mass index (BMI) was obtained from the physical examination and calculated as weight in kilograms (kg) divided by the square of height in meter. Weight was measured with an electronic scale, with a precision of 0.1 kg and maximum capacity of 150 kg, and height was measured with a portable stadiometer with a precision of 0.1m, respectively

STATISTICAL ANALYSIS

In order to calculate the data, Stata⁸ software was applied to perform descriptive and inferential statistical tests. Means, percentages and standard errors were estimated to describe the sample characteristics and the association between variables. Data research was analyzed by using two proportion test chi Square (X^2), however, if one expected value in the cell less than 5 (>20%) then fisher exact test was applied in accordingly.

Further, blood lead levels were left skewed and log transformed for analysis. Outcome variables, systolic and diastolic blood pressures were determined, respectively, inverse and log transformed to follow normal distribution. Multiple linear regression models were performed to examine relationships of blood lead with systolic and diastolic blood pressures comparing those participants in quartile 2 to 4 of blood lead levels with those in quartile 1. Multiple logistic regression analysis was used to evaluate the risk of hypertension also by categorizing blood lead in quartiles. This study also performed Pearson correlation analysis to verify the correlations between systolic and diastolic blood pressure with blood lead levels.

Regression models were constructed based on a priori knowledge and biologic association with blood pressure (age, sex, anti hypertensive medication use and blood lead log transformed). Other covariates were added to the model in two separated blocks: Model 1 – sex, age, race, income, education, antihypertensive medication and blood lead level; Statistical tests with p value < .05 were considered statistically significant.

RESULTS

The number of the selected sample was 35 fisherman who have been living and working in the

Makassar coastal areas. The characteristics of those respondents is presented in the table 1.

Table 1. Characteristics of respondent in the Makassar coastal area

Characteristics	Mean	Standard deviation	Minimum	Maximum
Age (years)	47.3	8.82	21	45
Sex (% male)	24.2	3.32	8.5	18.5
Working period (y)	22.3	4.51	1.2	11.2
Working duration(y)	18.2	8.56	11.9	26.9
Education (y)	5.2	9.78	9.20	17.3
IMT	11.2	4.71	10.0	32.0

The mean of age, sex, working period, working duration, education and IMT of the respondents were 47.3, 24.2, 22.3, 18.2, 5.2, and 11.2, respectively. The longest deviation standard was education with 9.78 whereas the shortest deviation standard was sex with

3.32. in addition the minimum SD was working period whereas the maximum was IMT with 32.0. Further, of the 35 respondents who had taken the blood sample and had blood measurement, the level of lead and hypertension information is described in the table 2.

Table 2. Result of respondents blood pressure measurement

Blood pressure status	Number	Percentage
Normal (140/90 mmHg)	8 person	22.9
Not normal ($\geq 140/90$ mmHg)	27 person	77.1
Number	35 Person	100

Hypertension defined as systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg or antihypertensive medication use.

Table 2 revealed that of the 35 respondents, there were 8 respondents who have a normal blood pressure and 27 have not normal blood pressure during the measurement. After the calculation, the lead (Pb) concentration in the blood with the incidence of hypertension can be found in the following table 3.

Table 3. Lead levels in the blood with the incidence of Hypertension both in pressure systolic and diastolic blood

Variables	Mean $\mu\text{gr/dL}$	Standard deviation	Minimum mmHg	Maximum mmHg
Lead in blood	27.6	17.56	2	89
Systolic blood pressure	144.6	13.32	98	123
Diastolic blood pressure	84.2	12.37	54	154

Table 3 indicated Pb levels in blood averaging 27.6 $\mu\text{gr/dL}$ with standard deviation 17.56 whereas the minimum value 2 and maximum value 89. In addition, the mean of systolic blood pressure 144.6 $\mu\text{gr/dL}$, standard deviation was 13.32, minimum value was 98 mmHg and maximum value 123 mmHg, mean diastolic blood pressure 84.2 $\mu\text{gr/dL}$, standard deviation 12.37,

and the minimum value 54 mmHg where maximum value of 154 mmHg.

DISCUSSION

This study indicated that there is a significant association between blood lead levels and the elevations in systolic and diastolic blood pressures. Result of

statistical test describe that there is a correlation between blood lead concentrations with hypertension incidence as well. The result showed that 27 respondents (77.1%) have high blood level, of those, there were 11 respondents suffer from hypertension, $p = 0,028$ smaller than $\alpha = 0,05$, it can be concluded there is significant correlation between Pb level in blood with hypertension incidence.

A study result also revealed that the average Pb of the water is about 22.00 $\mu\text{g}/\text{dl}$ although the levels are lower when compared to the safe level of Pb levels in the blood by WHO⁹ around 10-25 $\mu\text{g} / \text{dl}$. However, these values indicate that the average respondent has experienced lead poisoning in the blood because the levels have exceeded the poisoning limit of 5 $\mu\text{g} / \text{dl}$. Respondents who had high blood levels of Pb had greater risk of hypertension than those who had low blood levels of Pb. This result is consistent with Lippmann's assertion that the effect on blood pressure occurs at blood levels of Pb between 5-35 $\mu\text{g}/\text{dl}$. The study of Riyadina et al.¹⁰, which states that high blood levels of Pb have a relationship with the incidence of hypertension. Other studies suggested that there is sufficient evidence to infer a causal relationship of exposure to lead with hypertension, besides the modest strength observed in these associations^{11,12}. In addition, studies with general population also support that found positive associations between blood lead concentrations and elevation on both systolic and diastolic blood pressure^{13,14}. According to WHO⁹, Pb concentration in blood range between 10-30 $\mu\text{g}/\text{dl}$ for adults will lead to blood pressure systolic problem and eritrosit protoporphyrin. Lead in blood may generate some disturbance including hypertension.

It is differently to what has been investigated on the lead effects of hypertensive in studies involving workers exposed to high levels of lead, environmental factors exposure is often asymptomatic and as other toxic effects of lead. The magnitude of raised blood pressure depends on the magnitude of the exposure duration and frequency, the longer of exposure duration and frequency the higher magnitude of blood pressure^{15,16}. Another perspective on the health effects of lead-associated increases in blood pressure is to consider the effect of a unit increase in median BLL on the overall distribution of blood pressure values in a population and the effect on number of persons with blood pressure elevations associated with stroke and other endpoints. This was calculated by Schwartz¹⁷, who reported that a 1 $\mu\text{g}/\text{dL}$ increase in mean population blood lead in US

white males aged 40–59, distributed over the range of blood pressure values in this population, would increase the numbers of persons at risk for myocardial infarction by 3200 per year, of strokes by 1300 per year, as well as an overall increase in clinical hypertension of 635,000.

The inclusion of confounding factors in regression models has been considered very important to control for residuals in such analysis, especially when examining small effects as the influence of lead in blood pressure¹⁸. The inclusion of inappropriate covariates in regression models can attenuate associations between lead levels and the studied outcomes. Still, the absence of certain variables in the analysis may overestimate the strength of the association¹⁹. In this study, we included in the analysis risk factors for cardiovascular disease such as smoking, alcohol consumption, sex, age, socioeconomic and demographic variables and BMI to account for covariates that are known to be associated with blood pressure outcomes. Even so, some unmeasured variables not included in the analysis may influence the association of lead and blood pressure as, for example, pre-existing diseases, genetic aspects and renal function.

Furthermore, lead-induced oxidative stress reduces the availability of NO, which is an endogenous catalyst of several biochemical processes with an important role in cardiovascular system regulation, leading to endothelial dysfunction. Barbosa Jr et al.²⁰ reported that lead exposure exerted a significant inhibitory effect on NO production, as shown by analyses of nitrite concentrations in adults living in the Southeast region of Brazil, suggesting that this biological mechanism is possibly related to the increased cardiovascular risk associated with lead exposure.²⁰⁻²³

This study also included the measurement of multiple anthropometric and socioeconomic characteristics and the standardization of blood pressure measurements. However, some limitations of this study are the cross-section⁷ design that does not allow causal inferences about the association of blood lead and blood pressure; and the fact that we may not have measured all the potential confounders of the relationship between hypertension and lead exposure.

Differences observed among the results of other studies may be associated to the methodological design, such as study of bone lead measurement as a marker of lead exposure compared with blood lead and to stratified

statistical analysis which have shown that distinct risk factors for lead exposure exist among populations in Americans. Although family income and education level have been considered significant predictors of blood lead levels, the historically difference observed among income level in fact has been decreasing. Furthermore, some Pb-related outcomes, including blood pressure outcomes, may vary by race/ethnicity, providing some evidence that certain groups of people may be at higher risk of generating some various diseases at the end²⁴⁻²⁶.

CONCLUSIONS

There is a positive association between blood lead and diastolic blood pressure, and a significant association and hypertension in communities in the coastal area of Makassar aged 40 years or older. This results add knowledge about the health risks associated with current environmental exposures to lead in Makassar, and encourage efforts to prevent lead exposure in the general population.

Acknowledgement: We thanks to the Ministry of Education, the Republic of Indonesia for giving a fund for this research. Also to the Health laboratory staffs who have analyzed the accordingly

Conflict of Interests : We declare that there is no any potential conflict financial interests within this study and from the funding institution including the publication.

Ethics approval : All participants who participated on this research were asked to sign a consent form after the presentation of the research objectives in front of them. Every participant consented to the publication of data. At any interview, two copies of the consent form were issued, one being given to the participant another one kept by researcher.

REFERENCES

1. Toxicological profile for lead. [http://www.atsdr.cdc.gov/toxprofiles/tp13.pdf]. Accessed 12 July 2016.
2. U.S. Environmental Protection Agency. Air Quality Criteria for Lead. [https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=158823]. Accessed 12 July 2016.
3. Navas-Acien A, Guallar E, Silbergeld EK, Rothenberg SJ. Lead exposure and cardiovascular disease--a systematic review. *Environ Health Perspect.* 2007;115(3):472-82.
4. Schwartz BS, Hu H. Adult lead exposure: time for change. *Environ Health Perspect.* 2007;115(3):451-4.
5. WHO. Global status report on noncommunicable diseases 2010. Burden: mortality, morbidity and risk factors. 2010:9-31.
6. Paoliello MM, De Capitani EM. Environmental contamination and human exposure to lead in Brazil. *Rev Environ Contam Toxicol.* 2005;184:59-96.
7. Mallongi, A., P. Parkpian, P. Pataranawat, S. Chinwetkitvanich, 2015, Mercury distribution and its potential environmental and health risks in aquatic habitat at artisanal buladu gold mine in Gorontalo Province, Indonesia. *Pak. J. Nutr.* 14: 1010-1025.
8. STATA CORP. Stata Statistical Software: Release 14. StataCorp LP. 2015.
9. World Health Organization (WHO), 1997. Deteksi Dini Penyakit Akibat Kerja, Buku Kedokteran, Jakarta
10. Riyadina, W., M. Notosiswoyo, Anna M. Sirait, L. Tana, 2002, "Hubungan Antara Plumbum (Pb) dalam darah dengan hipertensi pada operator pompa bensin (SPBU)", *Buletin Penelitian Kesehatan*, 30 (2): 82-88.2006
11. Navas-Acien A, Guallar E, Silbergeld EK, Rothenberg SJ. Lead exposure and cardiovascular disease--a systematic review. *Environ Health Perspect.* 2007;115(3):472-82.
12. Glenn BS, Stewart WF, Links JM, Todd AC, Schwartz BS. The longitudinal association of lead with blood pressure. *Epidemiology.* 2003;14(1):30-6.
13. Bushnik T, Levallois P, D'Amour M, Anderson TJ, McAlister FA. Association between blood lead and blood pressure: Results from the Canadian Health Measures Survey (2007 to 2011). *Health Rep.* 2014;25(7):12-22.
14. Martin D, Glass TA, Bandeen-Roche K, Todd AC, Shi W, Schwartz BS. Association of blood lead and tibia lead with blood pressure and hypertension in a community sample of older adults. *Am J Epidemiol.* 2006;163(5):467-78.
15. Agency for Toxic Substances and Disease Registry.

- Toxicological Profile for Lead. [<http://www.atsdr.cdc.gov/toxprofiles/tp13.pdf>]. Accessed 12 July 2016.
16. U. S. (Environmental Protection Agency). Integrated Science Assessment for Lead. [<http://cfpub.epa.gov/ncea/cfm/recorddisplay.cfm?deid=255721-Download>]. Accessed 14 Oct 2015.
17. Schwartz J. Lead, blood pressure and cardiovascular disease in men. *Arch Environ Health*. 1995;50(1):31–7.
18. Nash D, Magder L, Lustberg M, Sherwin RW, Rubin RJ, Kaufmann RB, Silbergeld EK. Blood lead, blood pressure, and hypertension in perimenopausal and postmenopausal women. *Jama*. 2003;289(12):1523–32.
19. Martin D, Glass TA, Bandeen-Roche K, Todd AC, Shi W, Schwartz BS. Association of blood lead and tibia lead with blood pressure and hypertension in a community sample of older adults. *Am J Epidemiol*. 2006;163(5):467–78.
20. Barbosa Jr F, Sertorio JT, Gerlach RF, Tanus-Santos JE. Clinical evidence for lead-induced inhibition of nitric oxide formation. *Arch Toxicol*. 2006;80(12):811–6.
21. Rantetampang, A.L. and A. Mallongi, Risks Assessment Of Cadmium Through Aquatic Biota Consumption From Sentani Lake In Papua, Indonesia *International Journal of Engineering Research & Technology (IJERT)*, 2013. 2(6): p. 2493-2500. 2013.
22. Mallongi, A., Herawaty, 2015, Assessment of mercury accumulation in dry deposition, surface soil and rice grain in Luwuk gold mine, Central Sulawesi *Research Journal of Applied Sciences*, 10(1): 22-24, 2015.
23. Yi, Y., Z. Yang, and S. Zhang, Ecological risk assessment of heavy metals in sediment and human health risk assessment of heavy metals in fishes in the middle and lower reaches of the Yangtze River basin. *Environmental Pollution*, 2011. 159(10): p. 2575-2585. 2011.
24. U. S. (Environmental Protection Agency). Integrated Science Assessment for Lead. [<http://cfpub.epa.gov/ncea/cfm/recorddisplay.cfm?deid=255721-Download>]. Accessed 14 Oct 2015.
25. Scinicariello F, Yesupriya A, Chang MH, Fowler BA. Modification by ALAD of the association between blood lead and blood pressure in the U.S. population: results from the Third National Health and Nutrition Examination Survey. *Environ Health Perspect*. 2010;118(2):259–64.
26. Yang, Y., Comprehensive assessment of heavy metal contamination in sediment of the Pearl River Estuary and adjacent shelf. *Marine Pollution Bulletin*, 2012. 64(9): p. 1947-1955. 2012.

Paper 4

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

10%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1

repo.unand.ac.id

Internet Source

2%

2

Lei Han, Xiuxia Wang, Ruhui Han, Ming Xu, Yuan Zhao, Qianqian Gao, Huanxi Shen, Hengdong Zhang. "Association between blood lead level and blood pressure: An occupational population-based study in Jiangsu province, China", PLOS ONE, 2018

Publication

2%

3

www.ijsrp.org

Internet Source

2%

4

Submitted to Pennsylvania State System of Higher Education

Student Paper

1%

5

es.scribd.com

Internet Source

1%

6

Wen-Yi Yang, Jan A. Staessen. "Letter to editor: Blood pressure, hypertension and lead exposure", Environmental Health, 2018

Publication

1%

7	tamararubin.com Internet Source	1%
8	Ana Carolina Bertinde Almeida Lopes, Mariana Ragassi Urbano, André de Souza-Nogueira, Gustavo H. Oliveira-Paula et al. "Association of lead, cadmium and mercury with paraoxonase 1 activity and malondialdehyde in a general population in Southern Brazil", Environmental Research, 2017 Publication	1%
9	ijhs.org.sa Internet Source	1%
10	www.gssrr.org Internet Source	1%
11	cunyisph.org Internet Source	1%
12	ijt.arakmu.ac.ir Internet Source	1%
13	jamanetwork.com Internet Source	1%
14	vufind.katalog.k.utb.cz Internet Source	1%
15	eprints.ums.ac.id Internet Source	1%

Exclude quotes On

Exclude bibliography Off

Exclude matches < 1%